

The Bayes classifier

Theorem

The classifier $h^*(\mathbf{x}) := \arg \max_k \eta_k(\mathbf{x})$ satisfies

$$R(h^*) = \min R(h)$$

where the min is over all possible classifiers.

To calculate the Bayes classifier/Bayes risk, we need to know $\eta_k(\mathbf{x}) = \mathbb{P}[Y = k | X = \mathbf{x}]$

Alternatively, since $\pi_k f_{X|Y}(\mathbf{x}|k) = \eta_k(\mathbf{x}) f_X(\mathbf{x})$, to find the maximum $\eta_k(\mathbf{x})$ it is sufficient to know $\pi_k f_{X|Y}(\mathbf{x}|k)$

Linear discriminant analysis (LDA)

In linear discriminant analysis (LDA), we make the (strong) assumption that

$$X|Y = k \sim \mathcal{N}(\boldsymbol{\mu}_k, \boldsymbol{\Sigma})$$

for $k = 0, \dots, K - 1$

Here $\mathcal{N}(\boldsymbol{\mu}, \boldsymbol{\Sigma})$ is the multivariate Gaussian/normal distribution with mean $\boldsymbol{\mu}$ and covariance matrix $\boldsymbol{\Sigma}$

$$\phi(\mathbf{x}; \boldsymbol{\mu}, \boldsymbol{\Sigma}) = \frac{1}{(2\pi)^{d/2} |\boldsymbol{\Sigma}|^{1/2}} \exp \left\{ -\frac{1}{2} (\mathbf{x} - \boldsymbol{\mu})^T \boldsymbol{\Sigma}^{-1} (\mathbf{x} - \boldsymbol{\mu}) \right\}$$

Note: Each class has the same covariance matrix $\boldsymbol{\Sigma}$

Example

Suppose that $K = 2$

$$d_M^2(\mathbf{x}; \hat{\boldsymbol{\mu}}_0, \hat{\boldsymbol{\Sigma}}) - 2 \log \hat{\pi}_0 \stackrel{0}{\underset{1}{\leq}} d_M^2(\mathbf{x}; \hat{\boldsymbol{\mu}}_1, \hat{\boldsymbol{\Sigma}}) - 2 \log \hat{\pi}_1$$

It turns out that by setting

$$\mathbf{w} = \hat{\boldsymbol{\Sigma}}^{-1} (\hat{\boldsymbol{\mu}}_1 - \hat{\boldsymbol{\mu}}_0)$$

$$b = \frac{1}{2} \hat{\boldsymbol{\mu}}_0^T \hat{\boldsymbol{\Sigma}}^{-1} \hat{\boldsymbol{\mu}}_0 - \frac{1}{2} \hat{\boldsymbol{\mu}}_1^T \hat{\boldsymbol{\Sigma}}^{-1} \hat{\boldsymbol{\mu}}_1 + \log \frac{\hat{\pi}_1}{\hat{\pi}_0}$$

we can re-write this as

$$\mathbf{w}^T \mathbf{x} + b \stackrel{0}{\underset{1}{\leq}} 0 \quad \textit{linear classifier}$$

Challenges for LDA

The generative model is rarely valid

Moreover, the number of parameters to be estimated is

- class prior probabilities: $K - 1$
- means: Kd
- covariance matrix: $\frac{1}{2}d(d + 1)$

If d is small and n is large, then we can accurately estimate these parameters (provably, using Hoeffding)

If n is small and d is large, then we have more parameters than observations, and will likely obtain very poor estimates

- first apply a dimensionality reduction technique to reduce d (more on this later)
- assume a more structured covariance matrix

Another possible escape

Recall from the very beginning of the lecture that the Bayes classifier can be stated either in terms of maximizing $\pi_k f_{X|Y}(\mathbf{x}|k)$ or $\eta_k(\mathbf{x})$

In LDA, we are estimating $\pi_k f_{X|Y}(\mathbf{x}|k)$, which is equivalent to the full joint distribution of (X, Y)

All we **really** need is to be able to estimate $\eta_k(\mathbf{x})$

- we don't need to know $f_X(\mathbf{x})$

LDA commits one of the cardinal sins of machine learning:

***Never solve a more difficult problem
as an intermediate step***

Is there a better approach?

Another look at plugin methods

Suppose $K = 2$

$$\begin{aligned}\text{Define } \eta(\mathbf{x}) &= \eta_1(\mathbf{x}) \\ &= 1 - \eta_0(\mathbf{x})\end{aligned}$$

In this case, another way to express the Bayes classifier is as

$$h^*(\mathbf{x}) = \begin{cases} 1 & \text{if } \eta(\mathbf{x}) \geq 1/2 \\ 0 & \text{if } \eta(\mathbf{x}) < 1/2 \end{cases}$$

Note that we do not actually need to know the full distribution of (X, Y) to express the Bayes classifier

All we really need is to decide if $\eta(\mathbf{x}) \geq 1/2$

Gaussian case

Suppose that $K = 2$ and that $X|Y = k \sim \mathcal{N}(\mu_k, \Sigma)$

$$\begin{aligned}\eta(\mathbf{x}) &= \frac{\pi_1 \phi(\mathbf{x}; \mu_1, \Sigma)}{\pi_1 \phi(\mathbf{x}; \mu_1, \Sigma) + \pi_0 \phi(\mathbf{x}; \mu_0, \Sigma)} \\ &= \frac{\pi_1 e^{-\frac{1}{2}(\mathbf{x}-\mu_1)^T \Sigma^{-1}(\mathbf{x}-\mu_1)}}{\pi_1 e^{-\frac{1}{2}(\mathbf{x}-\mu_1)^T \Sigma^{-1}(\mathbf{x}-\mu_1)} + \pi_0 e^{-\frac{1}{2}(\mathbf{x}-\mu_0)^T \Sigma^{-1}(\mathbf{x}-\mu_0)}} \\ &= \frac{1}{1 + \frac{\pi_0}{\pi_1} e^{\frac{1}{2}(\mathbf{x}-\mu_1)^T \Sigma^{-1}(\mathbf{x}-\mu_1) - \frac{1}{2}(\mathbf{x}-\mu_0)^T \Sigma^{-1}(\mathbf{x}-\mu_0)}} \\ &= \frac{1}{1 + e^{-(\mathbf{w}^T \mathbf{x} + b)}}\end{aligned}$$

Logistic regression

This observation gives rise to another class of plugin methods, the most important of which is logistic regression, which implements the following strategy

1. Assume $\eta(\mathbf{x}) = \frac{1}{1 + e^{-(\mathbf{w}^T \mathbf{x} + b)}}$ ($\mathbf{w} \in \mathbb{R}^d, b \in \mathbb{R}$)

2. Directly estimate \mathbf{w}, b (somehow) from the data


3. Plug the estimate

$$\hat{\eta}(\mathbf{x}) = \frac{1}{1 + e^{-(\hat{\mathbf{w}}^T \mathbf{x} + \hat{b})}}$$

into the formula for the Bayes classifier

The logistic function

The function $\frac{1}{1+e^{-t}}$ is called a *logistic* function (or a *sigmoid* function in other contexts)



The logistic regression classifier

Denote the logistic regression classifier by

$$\hat{h}(\mathbf{x}) = 1_{\{\hat{\eta}(\mathbf{x}) \geq 1/2\}}(\mathbf{x})$$

Note that $\hat{h}(\mathbf{x}) = 1 \iff \hat{\eta}(\mathbf{x}) \geq \frac{1}{2}$


$$\iff \frac{1}{1 + \exp(-(\hat{\mathbf{w}}^T \mathbf{x} + \hat{b}))} \geq \frac{1}{2}$$

$$\iff \exp(-(\hat{\mathbf{w}}^T \mathbf{x} + \hat{b})) \leq 1$$

$$\iff (\hat{\mathbf{w}}^T \mathbf{x} + \hat{b}) \geq 0$$

So $\hat{h}(\mathbf{x}) = \begin{cases} 1 & \text{if } \hat{\mathbf{w}}^T \mathbf{x} + \hat{b} \geq 0 \\ 0 & \text{otherwise} \end{cases}$ *linear classifier*

Example



Estimating the parameters

Challenge: How to estimate the parameters for

$$\eta(\mathbf{x}) = \frac{1}{1 + e^{-(\mathbf{w}^T \mathbf{x} + b)}}$$

One possibility: $\mathbf{w} = \hat{\Sigma}^{-1} (\hat{\boldsymbol{\mu}}_1 - \hat{\boldsymbol{\mu}}_0)$

$$b = \frac{1}{2} \hat{\boldsymbol{\mu}}_0^T \hat{\Sigma}^{-1} \hat{\boldsymbol{\mu}}_0 - \frac{1}{2} \hat{\boldsymbol{\mu}}_1^T \hat{\Sigma}^{-1} \hat{\boldsymbol{\mu}}_1 + \log \frac{\hat{\pi}_1}{\hat{\pi}_0}$$

Alternative: *Maximum likelihood estimation*

For convenience, let's let $\boldsymbol{\theta} = (b, \mathbf{w})$

Note that $\eta(\mathbf{x})$ is really a function of both \mathbf{x} and $\boldsymbol{\theta}$, so we will use the notation $\eta(\mathbf{x}; \boldsymbol{\theta})$ to highlight this dependence

The *a posteriori* probability of our data

Suppose that we knew θ . Then we could compute

$$\begin{aligned}\mathbb{P}[y_i | \mathbf{x}_i; \theta] &= \mathbb{P}[Y_i = y_i | X_i = \mathbf{x}_i; \theta] \\ &= \begin{cases} \eta(\mathbf{x}_i; \theta) & \text{if } y_i = 1 \\ 1 - \eta(\mathbf{x}_i; \theta) & \text{if } y_i = 0 \end{cases} \\ &= \eta(\mathbf{x}_i; \theta)^{y_i} (1 - \eta(\mathbf{x}_i; \theta))^{1-y_i}\end{aligned}$$

Because of independence, we also have that

$$\begin{aligned}\mathbb{P}[y_1, \dots, y_n | \mathbf{x}_1, \dots, \mathbf{x}_n; \theta] &= \prod_{i=1}^n \mathbb{P}[y_i | \mathbf{x}_i; \theta] \\ &= \prod_{i=1}^n \eta(\mathbf{x}_i; \theta)^{y_i} (1 - \eta(\mathbf{x}_i; \theta))^{1-y_i}\end{aligned}$$

Maximum likelihood estimation

We don't actually know θ , but we do know y_1, \dots, y_n

Suppose we view y_1, \dots, y_n to be fixed, and view $\mathbb{P}[y_1, \dots, y_n | \mathbf{x}_1, \dots, \mathbf{x}_n; \theta]$ as just a function of θ

When we do this, $\mathcal{L}(\theta) = \mathbb{P}[y_1, \dots, y_n | \mathbf{x}_1, \dots, \mathbf{x}_n; \theta]$ is called the **likelihood** (or likelihood function)

The method of **maximum likelihood** aims to estimate θ by finding the θ that **maximizes** the **likelihood** $\mathcal{L}(\theta)$

In practice, it is often more convenient to focus on maximizing the **log-likelihood**, i.e., $\log \mathcal{L}(\theta)$

The log-likelihood

To see why, note that the likelihood in our case is given by

$$\mathcal{L}(\boldsymbol{\theta}) = \prod_{i=1}^n \eta(\mathbf{x}_i; \boldsymbol{\theta})^{y_i} (1 - \eta(\mathbf{x}_i; \boldsymbol{\theta}))^{1-y_i}$$

Thus, the log-likelihood is given by

$$\begin{aligned} \ell(\boldsymbol{\theta}) &= \log \mathcal{L}(\boldsymbol{\theta}) \\ &= \sum_{i=1}^n y_i \log \eta(\mathbf{x}_i; \boldsymbol{\theta}) + (1 - y_i) \log(1 - \eta(\mathbf{x}_i; \boldsymbol{\theta})) \end{aligned}$$

It is often easier to work with summations instead of products, and since the log is a monotonic transformation, maximizing $\ell(\boldsymbol{\theta})$ is equivalent to maximizing $\mathcal{L}(\boldsymbol{\theta})$

Simplifying the log-likelihood

Notation

- $\tilde{\mathbf{x}} = [1, x(1), \dots, x(d)]^T$
- $\boldsymbol{\theta} = [b, w(1), \dots, w(d)]^T$

This means that $\mathbf{w}^T \mathbf{x} + b = \boldsymbol{\theta}^T \tilde{\mathbf{x}}$, which lets us write

$$\eta(\mathbf{x}_i; \boldsymbol{\theta}) = \frac{1}{1 + e^{-\boldsymbol{\theta}^T \tilde{\mathbf{x}}_i}}$$

Thus, if we let $g(t) = \frac{1}{1+e^{-t}}$, then we can write

$$\ell(\boldsymbol{\theta}) = \sum_{i=1}^n y_i \log g(\boldsymbol{\theta}^T \tilde{\mathbf{x}}_i) + (1 - y_i) \log(1 - g(\boldsymbol{\theta}^T \tilde{\mathbf{x}}_i))$$

Simplifying the log-likelihood

Facts:

$$\log g(t) = \log \left(\frac{1}{1 + e^{-t}} \right) = -\log(1 + e^{-t})$$

$$\begin{aligned} \log(1 - g(t)) &= \log \left(1 - \frac{1}{1 + e^{-t}} \right) \\ &= \log \left(\frac{e^{-t}}{1 + e^{-t}} \right) = -t - \log(1 + e^{-t}) \\ &= \log \left(\frac{1}{1 + e^t} \right) = -\log(1 + e^t) \end{aligned}$$

Simplifying the log-likelihood

Facts:

$$\log g(t) = -\log(1 + e^{-t})$$

$$\log(1 - g(t)) = -t - \log(1 + e^{-t}) = -\log(1 + e^t)$$

Thus

$$\begin{aligned}\ell(\boldsymbol{\theta}) &= \sum_{i=1}^n y_i \log g(\boldsymbol{\theta}^T \tilde{\mathbf{x}}_i) + (1 - y_i) \log(1 - g(\boldsymbol{\theta}^T \tilde{\mathbf{x}}_i)) \\ &= \sum_{i=1}^n -y_i \log(1 + e^{-\boldsymbol{\theta}^T \tilde{\mathbf{x}}_i}) - \log(1 + e^{\boldsymbol{\theta}^T \tilde{\mathbf{x}}_i}) \\ &\quad - y_i(-\boldsymbol{\theta}^T \tilde{\mathbf{x}}_i - \log(1 + e^{-\boldsymbol{\theta}^T \tilde{\mathbf{x}}_i})) \\ &= \sum_{i=1}^n y_i \boldsymbol{\theta}^T \tilde{\mathbf{x}}_i - \log(1 + e^{\boldsymbol{\theta}^T \tilde{\mathbf{x}}_i})\end{aligned}$$

Maximizing the log-likelihood

How can we maximize

$$\ell(\boldsymbol{\theta}) = \sum_{i=1}^n y_i \boldsymbol{\theta}^T \tilde{\mathbf{x}}_i - \log(1 + e^{\boldsymbol{\theta}^T \tilde{\mathbf{x}}_i})$$

with respect to $\boldsymbol{\theta}$?

Find a $\boldsymbol{\theta}$ such that $\nabla \ell(\boldsymbol{\theta}) = \begin{bmatrix} \frac{\partial \ell(\boldsymbol{\theta})}{\partial \theta_1} \\ \vdots \\ \frac{\partial \ell(\boldsymbol{\theta})}{\partial \theta_{d+1}} \end{bmatrix} = 0$

(i.e., compute the partial derivatives and set them to zero)

Computing the gradient

It is not too hard to show that

$$\begin{aligned}\nabla \ell(\boldsymbol{\theta}) &= \sum_{i=1}^n \nabla \left(y_i \boldsymbol{\theta}^T \tilde{\mathbf{x}}_i - \log(1 + e^{\boldsymbol{\theta}^T \tilde{\mathbf{x}}_i}) \right) \\ &= \sum_{i=1}^n y_i \tilde{\mathbf{x}}_i - \tilde{\mathbf{x}}_i e^{\boldsymbol{\theta}^T \tilde{\mathbf{x}}_i} (1 + e^{\boldsymbol{\theta}^T \tilde{\mathbf{x}}_i})^{-1} \\ &= \sum_{i=1}^n \tilde{\mathbf{x}}_i (y_i - g(\boldsymbol{\theta}^T \tilde{\mathbf{x}}_i)) = 0\end{aligned}$$

This gives us $d + 1$ equations, but they are *nonlinear* and have no closed-form solution

Optimization

Throughout signal processing and machine learning, we will very often encounter problems of the form

$$\underset{\mathbf{x} \in \mathbb{R}^d}{\text{minimize}} f(\mathbf{x})$$

(or minimize $-\ell(\boldsymbol{\theta})$ for today)
 $\boldsymbol{\theta} \in \mathbb{R}^{d+1}$

In many (most?) cases, we cannot compute the solution simply by setting $\nabla f(\mathbf{x}) = 0$ and solving for \mathbf{x}

However, there are many powerful *algorithms* for finding \mathbf{x} using a computer

Gradient descent


A simple way to try to find the minimum of our objective function is to iteratively “*roll downhill*”

From \mathbf{x}^0 , take a step in the direction of the negative gradient

$$\mathbf{x}^1 = \mathbf{x}^0 - \alpha_0 \nabla f(\mathbf{x})|_{\mathbf{x}=\mathbf{x}^0} \quad \alpha_0 : \text{“step size”}$$

$$\mathbf{x}^2 = \mathbf{x}^1 - \alpha_1 \nabla f(\mathbf{x})|_{\mathbf{x}=\mathbf{x}^1}$$

⋮





Convergence of gradient descent

The core iteration of gradient descent is to compute

$$\mathbf{x}^{j+1} = \mathbf{x}^j - \alpha_j \nabla f(\mathbf{x})|_{\mathbf{x}=\mathbf{x}^j}$$


Note that if $\nabla f(\mathbf{x})|_{\mathbf{x}=\mathbf{x}^j} = 0$, then we have found the minimum and $\mathbf{x}^{j+1} = \mathbf{x}^j$, so the algorithm will terminate

If f is convex and sufficiently smooth, then gradient descent (with a fixed step size α) is guaranteed to converge to the global minimum of f




Step size matters!

Even though gradient descent provably converges, it can potentially take a while



Step size matters!

Even though gradient descent provably converges, it can potentially take a while



Newton's method

Also known as the Newton-Raphson method, this approach can be viewed as simply using the second derivative to automatically select an appropriate step size

$$\mathbf{x}^{j+1} = \mathbf{x}^j - (\nabla^2 f(\mathbf{x}))^{-1} \nabla f(\mathbf{x})|_{\mathbf{x}=\mathbf{x}^j}$$

**Hessian
matrix**

$$\nabla^2 f(\mathbf{x}) = \begin{bmatrix} \frac{\partial^2 f}{\partial x_1^2} & \frac{\partial^2 f}{\partial x_1 \partial x_2} & \cdots & \frac{\partial^2 f}{\partial x_1 \partial x_d} \\ \frac{\partial^2 f}{\partial x_2 \partial x_1} & \frac{\partial^2 f}{\partial x_2^2} & \cdots & \frac{\partial^2 f}{\partial x_2 \partial x_d} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial^2 f}{\partial x_d \partial x_1} & \frac{\partial^2 f}{\partial x_d \partial x_2} & \cdots & \frac{\partial^2 f}{\partial x_d^2} \end{bmatrix}$$

Optimization for logistic regression

The negative log-likelihood in logistic regression is a convex function

Both gradient descent and Newton's method are common strategies for setting the parameters in logistic regression

Newton's method is much faster when the dimension d is small, but is impractical when d is large

Why?

More on this on next week's homework

Comparison of plugin methods

Naïve Bayes, LDA, and logistic regression are all *plugin methods* that result in *linear* classifiers

Naïve Bayes

- plugin method based on density estimation
- scales well to high-dimensions and naturally handles mixture of discrete and continuous features

Linear discriminant analysis

- better if Gaussianity assumptions are valid

Logistic regression

- models only the distribution of $Y|X$, not (X, Y)
- valid for a larger class of distributions
- fewer parameters to estimate

Beyond plugin methods

Plugin methods can be useful in practice, but ultimately they are very limited

- There are always distributions where our assumptions are violated
- If our assumptions are wrong, the output is totally unpredictable
- Can be hard to verify whether our assumptions are right
- Require solving a more difficult problem as an intermediate step

For most of the remainder of this course will focus on *nonparametric* methods that avoid making such strong assumptions about the (unknown) process generating the data